

# Diastereotopic group selectivity and chemoselectivity of alkylidene carbene reactions on 8-oxabicyclo[3.2.1]oct-6-ene ring systems

Munro, Kevin R.; Male, Louise; Spencer, Neil; Grainger, Richard S.

DOI:  
[10.1039/c3ob41390j](https://doi.org/10.1039/c3ob41390j)

License:  
Creative Commons: Attribution-NonCommercial (CC BY-NC)

Document Version  
Publisher's PDF, also known as Version of record

Citation for published version (Harvard):  
Munro, KR, Male, L, Spencer, N & Grainger, RS 2013, 'Diastereotopic group selectivity and chemoselectivity of alkylidene carbene reactions on 8-oxabicyclo[3.2.1]oct-6-ene ring systems', *Organic and Biomolecular Chemistry*, vol. 11, no. 39, pp. 6856-6862. <https://doi.org/10.1039/c3ob41390j>

[Link to publication on Research at Birmingham portal](#)

**Publisher Rights Statement:**  
Eligibility for repository : checked 02/04/2014

## General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

Cite this: *Org. Biomol. Chem.*, 2013, **11**, 6856

## Diastereotopic group selectivity and chemoselectivity of alkylidene carbene reactions on 8-oxabicyclo[3.2.1]oct-6-ene ring systems†

Kevin R. Munro, Louise Male, Neil Spencer and Richard S. Grainger\*

$\alpha$ -Hydroxyalkylidene carbenes, generated from thermolysis of  $\alpha,\beta$ -epoxy-*N*-aziridinylimines, undergo diastereotopic group selective 1,5 C–H insertion reactions on 2,4-dimethyl-8-oxabicyclo[3.2.1]oct-6-ene ring systems. Protection of a tertiary alcohol at C-3 of the bridged oxabicyclo as a trimethylsilyl ether reverses the sense of diastereoselectivity. 1,5 C–H insertion into a methine adjacent to an OBn group, 1,5 O–R insertion into a tertiary alcohol (R = H) or silyl ether (R = TMS) at C-3 to form spirocyclic dihydrofurans, 1,2-rearrangement to an alkyne and fragmentation to a ketone are competing major pathways for 2-benzyloxy-substituted 8-oxabicyclo[3.2.1]oct-6-ene systems. Dihydrofuran formation is shown to be a result of substitution on the oxabicyclic ring system through comparison with other methods of alkylidene carbene formation.

Received 6th July 2013,  
Accepted 29th August 2013


DOI: 10.1039/c3ob41390j

[www.rsc.org/obc](http://www.rsc.org/obc)

### Introduction

The intramolecular 1,5-alkylidene carbene C–H insertion reaction is a powerful method for the synthesis of 5-membered, unsaturated carbo- and heterocyclic ring systems.<sup>1</sup> The ability to insert into unfunctionalized C–H bonds, with increasingly predictable chemo-, regio- and stereoselectivity, has led to widespread application in target synthesis.<sup>1,2</sup>

We are interested in the use of this reaction for the preparation of the A-ring of ingenol, a structurally complex diterpene of considerable synthetic and biological interest.<sup>3</sup> Our previous model studies established that 1,5 C–H insertion to give cyclopentenones **3** and **6** overrides the potentially competing 1,5 O–Si insertion pathway on structurally rigid 8-oxabicyclo[3.2.1]oct-6-ene ring systems **1** and **4** (Scheme 1).<sup>4,5</sup> However, these studies did not address the presence of an additional hydroxyl group within the cyclopentene A-ring of the natural product. Keen to avoid the need to introduce the required oxygenation at C-3 (ingenol numbering) post-annulation, we were attracted to the methodology of Kim for the generation of  $\alpha$ -hydroxyalkylidene carbenes **8** through thermolysis of  $\alpha,\beta$ -epoxy-*N*-aziridinylimines **7** (Scheme 2).<sup>6</sup> Depending upon substituents R<sup>1</sup>–R<sup>3</sup>, alkylidene carbenes **8** can undergo a



**Scheme 1** Previous chemo- and regioselective 1,5 C–H insertion of alkylidene carbenes on 8-oxabicyclo[3.2.1]oct-6-ene ring systems.


number of different reaction pathways,<sup>7,8</sup> with 1,5 C–H insertion leading to hydroxy-substituted cyclopentenones.<sup>6</sup>

In this paper we report our studies on the use of the Kim  $\alpha$ -hydroxyalkylidene carbene for hydroxycyclopentene annulation on two model 8-oxabicyclo[3.2.1]oct-6-ene ring systems.<sup>9</sup>

School of Chemistry, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK. E-mail: [r.s.grainger@bham.ac.uk](mailto:r.s.grainger@bham.ac.uk); Fax: +44 (0)121 4144403; Tel: +44 (0)121 4144465

†Electronic supplementary information (ESI) available: Experimental procedures and copies of <sup>1</sup>H and <sup>13</sup>C NMR for all new compounds. For ESI see DOI: 10.1039/c3ob41390j






**Scheme 2** Formation of  $\alpha$ -hydroxyalkylidene carbenes **8** from  $\alpha,\beta$ -epoxy-*N*-aziridinylimines **7**.

In addition to establishing the compatibility of this methodology on systems bearing oxygen functionality at C-3 of the oxabicyclo (a potential site of reactivity),<sup>5,7</sup> these studies were designed to probe two additional issues. The directing effect of the hydroxyl group in **8**, initially masked in the form of an epoxide **7**, is investigated through a formal diastereotopic group selective 1,5 C–H insertion reaction.<sup>10,11</sup> The influence of additional oxygenation, of relevance to the 5-hydroxyl group in ingenol, is investigated on C-2 benzyloxy-substituted oxabicycles. The formation of products arising from 1,5 O–Si and O–H insertion reactions on these latter systems are further probed using alternative methods of alkylidene carbene formation.

## Results and discussion

### Stereoselective synthesis of alkylidene carbene precursors from oxabicyclo[3.2.1]oct-6-en-3-ones

The Kim alkylidene carbene precursors,  $\alpha,\beta$ -epoxy-*N*-aziridinylimines **16a** ( $R = R' = \text{Me}$ ) and **16b** ( $R = \text{OBn}$ ,  $R' = \text{H}$ ), were prepared from the known 8-oxabicyclic ketones **9a**<sup>12</sup> and **9b**,<sup>13</sup> readily accessible from [4 + 3] cycloaddition reactions of substituted oxyallyl cations with furan (Scheme 3).<sup>14</sup> Stereoselective addition of the vinyl lithium species generated from iodoalkene **10**,<sup>15</sup> *syn* to the oxygen bridge, followed by removal of the TBS group *in situ*, gave the allylic diol **11**. Vanadium-catalysed epoxidation allowed for selective oxidation of the exocyclic alkene in **11a**.<sup>16</sup> In contrast, *m*-CPBA gave a mixture of products arising from additional reaction at the endocyclic alkene. The presence of a primary alcohol is also important to achieve the desired chemoselectivity – the corresponding TBS ether of **11a** (containing a tertiary alcohol) failed to undergo vanadium-catalysed epoxidation at either alkene. Vanadium-catalysed oxidation of allylic alcohol **11b** proceeded with 9 : 1 diastereoselectivity in favour of epoxide **12b**, the relative stereochemistry of which was confirmed on a later compound. The preferential formation of **12b** can be ascribed to epoxidation occurring through a conformation **11b'** where both the benzyl



**Scheme 3** Synthesis of alkylidene carbene precursors **16a** and **16b**.


ether and the primary alcohol can coordinate vanadium.<sup>17</sup> Further oxidation of the primary alcohol of **12** using Dess–Martin periodinane (DMP) in the presence of  $\text{NaHCO}_3$  gave the aldehyde **13**.<sup>18</sup> Protection of the tertiary alcohol as a TMS ether **14** followed by condensation with the hydrazine derived from ammonium salt **15**<sup>19</sup> gave the target *N*-aziridinylimine **16**.

### Diastereotopic group selective alkylidene carbene 1,5 C–H insertion reactions on 2,4-dimethyl-8-oxabicyclo[3.2.1]oct-6-enes

Heating a toluene solution of **16a** to reflux gave rise to a mixture of two diastereomeric cyclopentenols **18** and **19** in 64% overall yield, consistent with the generation and subsequent 1,5 C–H insertion of  $\alpha$ -hydroxyalkylidene carbene **17a** (Scheme 4). The two diastereoisomers **18** and **19** were separable by column chromatography, and their relative stereochemistry determined by nOe analysis. In the case of the minor isomer **19**, correlation is observed between the proton adjacent to the alcohol and the axial methine hydrogen on the tetrahydropyran ring, whereas no analogous signal is observed in the major isomer **18**.

Deprotection of the trimethylsilyl ether in **16a** with TBAF gave the tertiary alcohol **20a** in excellent yield (Scheme 4). The tertiary alcohol **20a** also gave rise to a mixture of cyclopentenols **22** and **23** upon heating to reflux in toluene, in lower overall yield (42%) and with the reversed sense of diastereoselectivity (1 : 2 ratio **22** : **23**) compared with the thermolysis of silyl ether **16a** to give cyclopentenols **18** and **19** (64%, 2 : 1 ratio **18** : **19**). Cyclopentenols **22** and **23** were isolable by column chromatography, and relative stereochemistry was again







**Scheme 4** Diastereotopic group selective 1,5 C–H insertion reactions of  $\alpha$ -hydroxyalkylidenes **17a** and **21a**.

determined by nOe correlation in the major diol **23**, with the structure of the minor diol **22** further confirmed by X-ray analysis (Fig. 1).<sup>†</sup>  $^1\text{H}$  NMR of the crude reaction mixture also showed the presence of dihydrofuran **24**, the product of 1,5 O–H insertion, and surprisingly the ketone **9a**, the product of C–C bond fragmentation. The ratio of C–H insertion products **22** and **23** to both **24** and **9a** was approximately 6 : 1 by integration of characteristic peaks in the  $^1\text{H}$  NMR of the crude reaction mixture.

The diastereoselectivity of the 1,5 C–H insertion reaction is clearly dependent on the C-3 substituent of the oxabicyclo, and while modest, is notably reversed for silyl ether **16a** and alcohol **20a**. The alcohol stereocentre in alkylidenes **17a** and **21a** renders the two potential bonds for 1,5 C–H insertion, C–H $^\alpha$  and C–H $^\beta$ , diastereotopic. The diastereotopic group selectivity<sup>20,21</sup> is controlled by the preferred orientation about



**Fig. 1** Crystal structure of hydroxycyclopentene **22** with ellipsoids drawn at the 50% probability level.



**Scheme 5** Stereochemical rationale for diastereotopic group selective C–H insertion.


the HO–C–C–OR carbon–carbon bond within the carbene (Scheme 5). For the TMS ether **17a**, insertion into C–H $^\beta$  is disfavoured compared to insertion into C–H $^\alpha$ , presumably due to unfavourable steric and/or electronic interactions between the silyl ether and the alcohol. For the tertiary alcohol **21a**, preferential insertion into C–H $^\beta$  can be rationalized by invoking a hydrogen bonding interaction, which fixes the conformation about the vicinal diol and orients the alkylidene carbene towards C–H $^\beta$ .

Attempts were made to increase the diastereoselectivity through functionalisation of the tertiary alcohol in **13a** with either a bulkier silyl group or an ester, which might alternatively act as a H-bond acceptor to the developing alcohol. Unfortunately the hindered nature of the tertiary alcohol in **13a** made it recalcitrant to reaction with TIPSOTf, TESOTf,  $\text{AcCl}$ ,  $\text{Cl}_2\text{CHCOCl}$  or  $\text{CF}_3\text{COCl}$ .

Further attempts to exploit an attractive interaction such as that proposed for **21a** (Scheme 5) instead promoted the unusual C–C bond fragmentation pathway leading to ketone **9a**. Running the thermolysis of **20a** in the presence of LiHMDS (to form a metal chelate) or phenylboronic acid (to form a cyclic boronate *in situ*<sup>22</sup>) gave **9a** in 22% and 52% isolated yield respectively, with no evidence of formation of 1,5 insertion products of an alkylidene carbene (Scheme 6). Alkoxy-silylcyclobutane **26**, expected to be Lewis acidic at Si through relief of ring strain,<sup>23</sup> was prepared in two steps from **13a** (yields

<sup>†</sup>**22**:  $\text{C}_{13}\text{H}_{18}\text{O}_3$ ,  $M = 222.27$ , orthorhombic,  $a = 13.4833(5)$ ,  $b = 11.3687(5)$ ,  $c = 14.6074(8)$  Å,  $U = 2239.13(18)$  Å<sup>3</sup>,  $T = 120(2)$  K, space group  $Pbca$ ,  $Z = 8$ , 24 957 reflections measured, 2558 unique ( $R_{\text{int}} = 0.0685$ ) which were used in all calculations. The final  $R_1$  was 0.0494 ( $I > 2\sigma(I)$ ) and  $wR(F_2)$  was 0.1200 (all data). CCDC 946066 contains the supplementary crystallographic data for this paper.





**Scheme 6** Fragmentation to ketone **9a**.


unoptimized). Again refluxing **26** in toluene resulted in fragmentation to form the ketone **9a**. Although the mechanism of this fragmentation has not been elucidated, a common factor may be the possible impedance of the proton transfer step in alkylidene carbene formation (Scheme 2), allowing alternative pathways to dominate.<sup>24</sup> This unusual pathway was further demonstrated to be a function of the aziridinyliimine – the epoxy aldehyde **25** proved stable to refluxing toluene, and could be recovered in good yield.

#### Chemoselectivity in $\alpha$ -hydroxyalkylidene carbene insertion reactions on 2-benzyloxy-8-oxabicyclo[3.2.1]oct-6-enes

Thermolysis of the TMS ether **16b** gave a relatively complex mixture of products, from which hydroxycyclopentene **27**, the product of 1,5 C–H insertion of alkylidene carbene **17b** adjacent to oxygen, could be isolated in 41% yield (Scheme 7).

nOe analysis of **27** showed a correlation between the proton adjacent to the hydroxyl group and one of the benzylic hydrogens, thus establishing the stereochemistry of major epoxide **12b** in the synthetic route to **16b** (Scheme 3). <sup>1</sup>H NMR analysis of the crude reaction mixture showed **27** to be the major product, obtained in an approximately 2.4 : 1 : 1.2 ratio along with silyldihydrofuran **28**, the product of 1,5 O–Si insertion, and the ketone **9b**, the product of C–C bond fragmentation. The structure of **28** within the reaction mixture was assigned on the basis of comparison with related tetrahydrofurans (*vide infra*), however it could not be separated by column chromatography. The combined yield of **27**, **28** and **9b** was 79%.

Removal of the TMS group from **16b** with TBAF gave the tertiary alcohol **20b**, which could not be adequately purified by column chromatography and so was instead taken directly into the thermolysis reaction to form the  $\alpha$ -hydroxyalkylidene carbene **21b**. Heating a toluene solution of the crude tertiary alcohol **20b** at reflux also gave rise to a relatively complex mixture of products, which could, however, be separated by column chromatography. Hydroxycyclopentene **29**, arising from 1,5 C–H insertion next to oxygen, and dihydrofuran **30**,



**Scheme 7** Regio- and chemoselectivity in alkylidene carbene insertion on 2-benzyloxy-substituted 8-oxabicyclo[3.2.1]oct-6-ene ring systems.

the product of O–H insertion, were the two major compounds, formed in an approximately 1 : 1 ratio. Significant amounts of ketone **9b** were again formed, and it was also possible to isolate small quantities of the alkyne **31**, the product of 1,2-rearrangement of the intermediate alkylidene carbene.

Alkylidene carbenes show a general preference for insertion into more substituted C–H bonds, tertiary > secondary > primary, a trend which is rationalized by the interaction of the more electron-rich C–H bond with the electron-deficient carbene.<sup>25</sup> The preferential 1,5-insertion of alkylidene carbenes into C–H bonds adjacent to heteroatoms is also well established in the literature. A recent comprehensive study by Lee has shown there is a directionality to the activating effect of an oxygen on an adjacent C–H bond. An ether oxygen can activate an adjacent C–H bond towards insertion through lone-pair donation into the  $\sigma_{C-H}^*$  antibonding orbital when appropriately aligned.<sup>2</sup> For both **17b** and **21b** 1,5 C–H insertion is completely selective for insertion into the methine group adjacent to the benzylic oxygen over the alternative 1,5 C–H insertion into the methylene group. The electronic and






stereoelectronic activation of the tertiary C–H apparently overrides, or acts in concert with, any potential directing effect of the alcohol stereocentre in **17b** and **21b**.

More surprising is the increased predominance of additional reaction pathways observed for the 2-benzyloxy-substituted oxabicyclo[3.2.1]oct-6-ene ring systems **16b** and **20b** compared to the 2,4-dimethyl-substituted systems **16a** and **20a**. Analogous O–Si/O–H insertion, fragmentation to ketone **9a**, and 1,2-migration are all pathways available to **16a** and **20a**, yet only occurred to a minor extent, or not at all. 1,2-Alkyl migration in particular is rarely observed for dialkyl-substituted alkylidene carbenes.<sup>10e,26,27</sup> Either 1,5 C–H insertion is slower than might be expected based on the additional stereoelectronic activation by the OBn group compared to the methyl groups in **17a** and **21a**, or structural features in the benzyloxy-substituted system promote these competing reaction pathways.

### Comparison with other methods of alkylidene carbene generation

We do not have a satisfactory rationale for this divergence in reactivity with substitution pattern, but the formation of significant quantities of enol ethers **28** and **30** is perhaps not as surprising as the fact that products arising from formal 1,5 O–Si were not observed in the thermolysis of **16a**, and the formation of dihydrofuran **24** from O–H insertion is only a minor pathway in the thermolysis **20a** (Scheme 4). Previous studies have shown that 1,5 O–Si and O–H insertion generally predominates over 1,5 C–H insertion where both pathways can operate.<sup>7,28</sup>

We have previously<sup>5</sup> ascribed the lack of O–Si insertion in the alkyl-substituted oxabicycles **1** and **4** (Scheme 1) to the preferred geometry about the C–OSiMe<sub>3</sub> bond, which orients the lone pairs on oxygen away from the alkylidene carbene (represented as the free carbenes **2** and **5**, but better envisioned as metal carbenoids<sup>29</sup>). Why then should O–Si and O–H insertion be favoured to such an extent for the benzyloxy-substituted systems? While the nature of the alkylidene carbene is different to that shown in Scheme 1, and the higher temperature at which it is being generated should increase bond rotation and may allow these alternative pathways to operate, this does not explain the lack of O–Si insertion in **16a**, and the relatively small amount of O–H insertion observed for **20a**. In order to separate any potential role of the developing hydroxyl group in directing the product outcome, *e.g.* through hydrogen bonding with the benzylic oxygen which may orientate the carbene away from the C–H insertion site, we have extended our prior studies in systems lacking this feature. In this way we hoped to determine the relative importance of the structure of the oxabicyclic *vs.* the nature of the carbene generation method on the product outcome.<sup>30</sup> Two classical methods of carbene generation were chosen, the base-mediated  $\alpha$ -elimination of a chloroalkene, and the reaction of a ketone with lithiated trimethylsilyldiazomethane, which generates the carbene through sequential Peterson olefination and loss of nitrogen.<sup>2,31</sup>



**Scheme 8** Alternative methods of alkylidene carbene generation.

$\beta$ -Silyloxyketone **34** was prepared in three steps from 2,4-dimethyl-8-oxabicyclo[3.2.1]oct-6-ene (**9a**) as previously described (Scheme 8).<sup>5</sup> Treatment of ketone **34** with lithium (trimethylsilyl)diazomethane gave cyclopentene **3**, the same product of C–H insertion we observed when generating the carbene from deprotonation of chloroalkene **1** (Scheme 1). Hence all three carbene generation methods on the 2,4-dimethyl-3-trimethylsilyloxy-substituted oxabicyclic systems give products of 1,5 C–H insertion, with no evidence of significant dihydrofuran formation through 1,5 O–Si insertion.

For the 2-benzyloxy-substituted system we employed a route previously developed for the synthesis of chloroalkene **4** (Scheme 1).<sup>5</sup> Addition of the dianion of methylacetoacetate to ketone **9b** gave  $\beta$ -ketoester **35** in good yield. Protection of the tertiary alcohol as a trimethylsilyl ether followed by Krapcho decarboxylation gave the  $\beta$ -silyloxyketone **36**. Subsequent Wittig reaction gave chloroalkene **37** as a 1:1 mixture of double bond isomers (Scheme 8).



The alkylidene carbenes generated from both **36** and **37** gave a mixture of compounds arising from both 1,5 C–H insertion and 1,5 O–Si insertion, which were separable after treatment of the crude reaction mixture with TBAF to liberate the tertiary alcohol **39**. The ratio of **38**:**39** changes with the method of carbene generation. Using lithium (trimethylsilyl)diazomethane O–Si insertion to give **38** is the dominant reaction pathway, whereas cyclopentene **39** is slightly favoured through deprotonation of chloroalkene **37**. Although there is a difference in product ratio with the method of generation of the carbene,<sup>30</sup> these results, and those using the Kim alkylidene carbene precursor, show a clear overall trend. The presence of a benzyloxy-substituent at C-2 of the oxabicyclic ring system controls the regioselectivity of 1,5 C–H insertion, but also leads to significant quantities of 1,5 O–Si insertion, irrespective of the method of carbene generation. In neither case was fragmentation to ketone **9b** nor 1,2-rearrangement to an alkyne observed, suggesting these pathways can be ascribed to the nature of the  $\alpha$ -hydroxyalkylidene carbene, or the conditions under which it is generated.

## Conclusions

Stereocontrolled annulation of hydroxycyclopentenones onto 8-oxabicyclo[3.2.1]oct-6-en-3-ones can be achieved using Kim's alkylidene carbene methodology, thus increasing the utility of these readily accessible and synthetically versatile ring systems.<sup>9,32</sup> A benzyloxy group at C-2 of the oxabicycle controls the position 1,5 C–H insertion, but results in increased amounts of products arising from alternative reaction pathways, compared with 2,4-dimethyl-substituted systems. Formation of spirocyclic dihydrofurans in 2-benzyloxy-substituted oxabicycles occurs irrespective of the type of alkylidene carbene employed, and is the dominant reaction pathway using lithium trimethylsilyldiazomethane. Additional reaction pathways, including a previously unobserved C–C bond fragmentation, occur upon heating  $\alpha,\beta$ -epoxy-*N*-aziridinylimines in refluxing toluene, compared to systems where the alkylidene carbene is generated at lower temperatures, and which lack an  $\alpha$ -hydroxyl group.

## Acknowledgements

We thank the EPSRC and University of Birmingham for funding. We thank the EPSRC UK National Crystallography Service at the University of Southampton for the collection of the crystallographic data.<sup>33</sup> The NMR spectrometers used in this research were obtained through Birmingham Science City: Innovative Uses for Advanced Materials in the Modern World (West Midlands Centre for Advanced Materials Project 2), with support from Advantage West Midlands (AWM) and part funded by the European Regional Development Fund (ERDF).

## Notes and references

- For reviews on alkylidene carbenes see: W. Kirmse, *Angew. Chem., Int. Ed. Engl.*, 1997, **36**, 1164; R. Knorr, *Chem. Rev.*, 2004, **104**, 3795.
- For lead references see: J.-C. Zheng, S. Y. Yun, C. Sun, N.-K. Lee and D. Lee, *J. Org. Chem.*, 2011, **76**, 1086.
- For reviews on the total synthesis of ingenol see: I. Kuwajima and K. Tanino, *Chem. Rev.*, 2005, **105**, 4661; J. K. Cha and O. L. Epstein, *Tetrahedron*, 2006, **62**, 1329; T. Busch and A. Kirschning, *Nat. Prod. Rep.*, 2008, **25**, 318.
- R. S. Grainger, R. B. Owoare, P. Tisselli and J. W. Steed, *J. Org. Chem.*, 2003, **68**, 7899.
- R. S. Grainger and R. B. Owoare, *Org. Lett.*, 2004, **6**, 2961.
- S. Kim and C. M. Cho, *Tetrahedron Lett.*, 1994, **35**, 8405.
- S. Kim and C. M. Cho, *Tetrahedron Lett.*, 1995, **36**, 4845.
- H.-Y. Lee and Y. Kim, *J. Am. Chem. Soc.*, 2003, **125**, 10156.
- For reviews on the synthetic utility of 8-oxabicyclo[3.2.1]oct-6-en-3-one ring systems see: P. Chiu and M. Lautens, in *Topics in Current Chemistry*, ed. P. Metz, Springer-Verlag, New York, 1997, vol. 190, pp. 1–85; I. V. Hartung and H. M. R. Hoffmann, *Angew. Chem., Int. Ed.*, 2004, **43**, 1934.
- For previous examples of regio- and diastereoselective alkylidene carbene 1,5 C–H insertion reactions controlled by an existing stereocentre in the connecting chain see: (a) D. F. Taber and R. P. Meagley, *Tetrahedron Lett.*, 1994, **35**, 7909; (b) S. Ohira, T. Sawamoto and M. Yamato, *Tetrahedron Lett.*, 1995, **36**, 1537; (c) D. F. Taber and H. Yu, *J. Org. Chem.*, 1997, **62**, 1687; (d) D. F. Taber, T. E. Christos, T. D. Neubert and D. Batra, *J. Org. Chem.*, 1999, **64**, 9673; (e) L. F. Walker, A. Bourghida, S. Connolly and M. Wills, *J. Chem. Soc., Perkin Trans. 1*, 2002, 965; (f) G. Hobley, K. Stuttle and M. Wills, *Tetrahedron*, 2003, **59**, 4739; (g) A. N. Van Nhien, E. Soriano, J. Marco-Contelles and D. Postel, *Carbohydr. Res.*, 2009, **344**, 1605; (h) A. N. Van Nhien, R. Cordonnier, M.-D. Le Bas, S. Delacroix, E. Soriano, J. Marco-Contelles and D. Postel, *Tetrahedron*, 2009, **65**, 9378; (i) S. Lee and H.-Y. Lee, *Bull. Korean Chem. Soc.*, 2010, **31**, 557.
- For desymmetrization of 8-oxabicyclo[3.2.1]octane ring systems through metal-catalysed asymmetric intramolecular C–H insertion see: P. Chiu, X. Zhang and R. Y. Y. Ko, *Tetrahedron Lett.*, 2004, **45**, 1531; X. Zhang, Z. Li, J. C. K. Chu and P. Chiu, *Tetrahedron Lett.*, 2011, **52**, 6763.
- M. Lautens and G. Bouchain, *Org. Synth.*, 2002, **79**, 251.
- M. Vidal-Pascual, C. Martínez-Lamenca and H. M. R. Hoffmann, *Org. Synth.*, 2006, **83**, 61.
- For recent reviews on the [4 + 3] cycloaddition reaction see: M. Harmata, *Chem. Commun.*, 2010, **46**, 8886; M. Harmata, *Chem. Commun.*, 2010, **46**, 8904; A. G. Lohse and R. P. Hsung, *Chem.-Eur. J.*, 2011, **17**, 3812.
- R. Baker and J. L. Castro, *J. Chem. Soc., Perkin Trans. 1*, 1990, 47; G. R. Scarlato, J. A. DeMattei, L. S. Chong, A. K. Ogawa, M. R. Lin and R. W. Armstrong, *J. Org. Chem.*, 1996, **61**, 6139; T. Motozaki, K. Sawamura, A. Suzuki,



- K. Yoshida, T. Ueki, A. Ohara, R. Munakata, K.-i. Takao and K.-i. Tadano, *Org. Lett.*, 2005, **7**, 2261.
- 16 K. B. Sharpless and R. C. Michaelson, *J. Am. Chem. Soc.*, 1973, **95**, 6136.
- 17 M. Kitamura, M. Isobe, Y. Ichikawa and T. Goto, *J. Org. Chem.*, 1984, **49**, 3517.
- 18 A. Fürstner, O. Larionov and S. Flügge, *Angew. Chem., Int. Ed.*, 2007, **46**, 5545.
- 19 R. K. Müller, R. Joos, D. Felix, J. Schreiber, C. Wintner and A. Eschenmoser, *Org. Synth.*, 1976, **55**, 114.
- 20 For reviews on diastereotopic group selective reactions see: R. W. Hoffmann, *Synthesis*, 2004, 2075; P. Herrmann and T. Bach, *Chem. Soc. Rev.*, 2011, **40**, 2022.
- 21 For previous contributions from our group in the area of diastereotopic group selective reactions see: R. S. Grainger, P. Tisselli and J. W. Steed, *Org. Biomol. Chem.*, 2004, **2**, 151; T. C. Casey, J. Carlisle, P. Tisselli, L. Male, N. Spencer and R. S. Grainger, *J. Org. Chem.*, 2010, **75**, 7461.
- 22 P. J. Duggan and E. M. Tyndall, *J. Chem. Soc., Perkin Trans. 1*, 2002, 1325.
- 23 A. G. Myers, S. E. Kephart and H. Chen, *J. Am. Chem. Soc.*, 1992, **114**, 7922; S. E. Denmark, B. D. Griedel, D. M. Coe and M. E. Schnute, *J. Am. Chem. Soc.*, 1994, **116**, 7026; K. Matsumoto, K. Oshima and K. Utimoto, *J. Org. Chem.*, 1994, **59**, 7152.
- 24 For a review on the chemistry of *N*-aziridinylimines see: W. Kirmse, *Eur. J. Org. Chem.*, 1998, 201.
- 25 J. Wolinsky, G. W. Clark and P. C. Thorstenson, *J. Org. Chem.*, 1976, **41**, 745; J. C. Gilbert, D. H. Giamalva and U. Weerasooriya, *J. Org. Chem.*, 1983, **48**, 5251.
- 26 D. J. Wardrop and E. G. Bowen, *Chem. Commun.*, 2005, 5106.
- 27 Alkyne **30** can arise through formal migration of either a methyl group or a hydroxyalkyl group. Wills has discussed the relative migratory aptitude of methyl vs. isopropyl vs. alkoxy towards alkylidene carbenes: ref. 10e.
- 28 K. Miwa, T. Aoyama and T. Shioiri, *Synlett*, 1994, 461; K. S. Feldman and M. L. Wroblewski, *J. Org. Chem.*, 2000, **65**, 8659; H.-J. Gais, L. R. Reddy, G. S. Babu and G. Raabe, *J. Am. Chem. Soc.*, 2004, **126**, 4859. For an exception see ref. 10f.
- 29 M. Braun, *Angew. Chem., Int. Ed.*, 1998, **37**, 430.
- 30 For an example of where the method of generation of the alkylidene carbene changes the regioselectivity of 1,5 C-H insertion see: D. F. Taber and T. E. Christos, *Tetrahedron Lett.*, 1997, **38**, 4927.
- 31 S. Ohira, K. Okai and T. Moritani, *J. Chem. Soc., Chem. Commun.*, 1992, 721; D. J. Wardrop, W. M. Zhang and J. Fritz, *Org. Lett.*, 2002, **4**, 489; C. J. Hayes, D. A. Bradley and N. A. Thomson, *J. Org. Chem.*, 2006, **71**, 2661.
- 32 For representative methodology and application of cyclopentane annulation on 8-oxabicyclo[3.2.1]octan-3-one ring systems see: Á. M. Montaña and K. M. Nicholas, *J. Org. Chem.*, 1990, **55**, 1569; P. A. Wender and F. E. McDonald, *J. Am. Chem. Soc.*, 1990, **112**, 4956; M. Nowakowski and H. M. R. Hoffmann, *Tetrahedron*, 1997, **53**, 4331; G. A. Molander and S. C. Jeffrey, *Tetrahedron Lett.*, 2002, **43**, 359; G. A. Molander, B. Czako and D. J. St. Jean Jr., *J. Org. Chem.*, 2006, **71**, 1172; Á. M. Montaña, S. Ponzano, G. Kociok-Köhn, M. Font-Bardia and X. Solans, *Eur. J. Org. Chem.*, 2007, 4383; J. Xu, E. J. E. Caro-Diaz and E. A. Theodorakis, *Org. Lett.*, 2010, **12**, 3708; Á. M. Montaña, S. Ponzano, C. Batalla and M. Font-Bardia, *Tetrahedron*, 2012, **68**, 8276; J. Wang, S.-G. Chen, B.-F. Sun, G.-Q. Lin and Y.-J. Shang, *Chem.-Eur. J.*, 2013, **19**, 2539.
- 33 S. J. Coles and P. A. Gale, *Chem. Sci.*, 2012, **3**, 683.

